

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

NASA TECHNICAL MEMORANDUM

NASA TM 78156

(NASA-TM-78156) EQUAL PERIOD ORBIT
DETERMINATION FOR THE SOLAR SYSTEM PLANETS
(NASA) 37 p HC A03/MF A01 CSCL 22A

N78-18092

Unclas

G3/13 05967

EQUAL PERIOD ORBIT DETERMINATION FOR THE SOLAR SYSTEM PLANETS

By Wilbur H. Funston
Data Systems Laboratory

January 10, 1978

NASA



*George C. Marshall Space Flight Center
Marshall Space Flight Center, Alabama*

1. REPORT NO. NASA TM 78156		2. GOVERNMENT ACCESSION NO.		3. RECIPIENT'S CATALOG NO.	
4. TITLE AND SUBTITLE Equal Period Orbit Determination for the Solar System Planets				5. REPORT DATE January 10, 1978	
				6. PERFORMING ORGANIZATION CODE	
7. AUTHOR(S) Wilbur H. Funston				8. PERFORMING ORGANIZATION REPORT #	
9. PERFORMING ORGANIZATION NAME AND ADDRESS George C. Marshall Space Flight Center Marshall Space Flight Center, Alabama 35812				10. WORK UNIT NO.	
				11. CONTRACT OR GRANT NO.	
12. SPONSORING AGENCY NAME AND ADDRESS National Aeronautics and Space Administration Washington, D.C. 20546				13. TYPE OF REPORT & PERIOD COVERED Technical Memorandum	
				14. SPONSORING AGENCY CODE	
15. SUPPLEMENTARY NOTES Prepared by Data Systems Laboratory, Science and Engineering.					
16. ABSTRACT <p>Rendezvous and docking maneuvers by spacecraft often make use of equi-period orbits. For example, a spacecraft in circular orbit about the Earth's Moon may put an observation module into an elliptical orbit which lowers the perigee for a closer observation of the lunar surface. Use of equi-period orbits will bring the observation module back to rendezvous with the other module which has remained in circular orbit. There will be one rendezvous opportunity each revolution.</p> <p>This report presents the determination of a set of equations for equi-period orbits work and a set of figures for quick reference.</p>					
17. KEY WORDS			18. DISTRIBUTION STATEMENT Unclassified — Unlimited		
19. SECURITY CLASSIF. (of this report) Unclassified		20. SECURITY CLASSIF. (of this page) Unclassified		21. NO. OF PAGES 37	
				22. PRICE NTIS	

TABLE OF CONTENTS

	Page
INTRODUCTION	1
DEVELOPMENT OF EQUATIONS FOR EQUI-PERIOD ORBITS	22
APPENDIX — VALUES USED FOR TABLES	30
BIBLIOGRAPHY	31

LIST OF ILLUSTRATIONS

Figure	Title	Page
1.	Velocity angle change for Earth	2
2.	Velocity angle change for Earth's Moon	3
3.	Velocity angle change for Mercury	4
4.	Velocity angle change for Venus	5
5.	Velocity angle change for Mars	6
6.	Velocity angle change for Jupiter	7
7.	Velocity angle change for Saturn	8
8.	Velocity angle change for Uranus	9
9.	Velocity angle change for Neptune	10
10.	Velocity angle change for Pluto	11
11.	Velocity change for Earth	12
12.	Velocity change for Earth's Moon	13
13.	Velocity change for Mercury	14
14.	Velocity change for Venus	15
15.	Velocity change for Mars	16
16.	Velocity change for Jupiter	17
17.	Velocity change for Saturn	18
18.	Velocity change for Uranus	19
19.	Velocity change for Neptune	20

LIST OF ILLUSTRATIONS (Concluded)

Figure	Title	Page
20.	Velocity change for Pluto	21
21.	Two-body orbital configuration	22
22.	Orbital vectors	25

TECHNICAL MEMORANDUM

EQUAL PERIOD ORBIT DETERMINATION FOR THE SOLAR SYSTEM PLANETS

INTRODUCTION

Rendezvous and docking maneuvers by spacecraft often make use of equi-period orbits. For example, a spacecraft in circular orbit about the Earth's Moon may put an observation module into an elliptical orbit which lowers the perigee for a closer observation of the lunar surface. Use of equi-period orbits will bring the observation module back to rendezvous with the other module which has remained in circular orbit. There will be one rendezvous opportunity each revolution.

Only the application of an impulse to the spacecraft in the proper attitude is required to change between circular and elliptical equi-period orbits. The orbit transfer is accomplished by applying sufficient impulse to change the velocity by:

$$dv = R_o \sqrt{2g_o} \left[\frac{1}{R_o + h} \left(1 - \left[\frac{(R_o + h)^2 - (h - h_p)^2}{(R_o + h)^2} \right]^{1/2} \right) \right]^{1/2} \quad (1)$$

The equi-period orbit velocity vector will be at an angle θ degrees from the circular velocity:

$$\theta = \arctan \frac{\frac{h}{R_o} - \frac{h_p}{R_o}}{\left[1 + \frac{2h}{R_o} + \frac{2hh_p}{R_o^2} - \frac{h^2}{R_o^2} \right]^{1/2}} \quad (2)$$

The equations in this report are straightforward and are used to generate Figures 1 through 20. The constants for these calculations are presented in the Appendix. All figures are for the drag free case and are duly noted.




Figure 1. Velocity angle change (θ) for Earth.

ORIGINAL PAGE IS
OF POOR QUALITY




Figure 3. Velocity angle change (θ) for Mercury.

ORIGINAL PAGE IS
OF POOR QUALITY




Figure 4. Velocity angle change (θ) for Venus.

ORIGINAL PAGE IS
OF POOR QUALITY




Figure 5. Velocity angle change (θ) for Mars.

VELOCITY ANGLE CHANGE VERSUS ORBIT ALTITUDE FOR EQUAL PERIOD CIRCULAR AND ELLIPTICAL ORBITS ABOUT SATURN




Figure 7. Velocity angle change (θ) for Saturn.




Figure 9. Velocity angle change (θ) for Neptune.




Figure 10. Velocity angle change (θ) for Pluto.

ORIGINAL PAGE IS
OF POOR QUALITY

VELOCITY CHANGE VERSUS ORBIT ALTITUDE FOR EQUAL PERIOD CIRCULAR AND ELLIPTICAL ORBITS ABOUT EARTH




Figure 11. Velocity change (ΔV) for Earth.

VELOCITY CHANGE VERSUS ORBIT ALTITUDE FOR EQUAL
PERIOD CIRCULAR AND ELLIPTICAL ORBITS ABOUT EARTH'S MOON




Figure 12. Velocity change (ΔV) for Earth's Moon.

ORIGINAL PAGE IS
OF POOR QUALITY

VELOCITY CHANGE VERSUS ORBIT ALTITUDE FOR EQUJAL PERIOD
CIRCULAR AND ELLIPTICAL ORBITS ABOUT MERCURY




Figure 13. Velocity change (ΔV) for Mercury.




Figure 14. Velocity change (ΔV) for Venus.

CIRCULAR ORBIT ALTITUDE - H (km)

Figure 15. Velocity change (ΔV) for Mars.

VELOCITY CHANGE VERSUS ORBIT ALTITUDE FOR EQUAL PERIOD CIRCULAR AND ELLIPTICAL ORBITS ABOUT SATURN




Figure 17. Velocity change (ΔV) for Saturn.

VELOCITY CHANGE VERSUS ORBIT ALTITUDE FOR EQUAL PERIOD
CIRCULAR AND ELLIPTICAL ORBITS ABOUT URANUS




Figure 18. Velocity change (ΔV) for Uranus.

VELOCITY CHANGE VERSUS ORBIT ALTITUDE FOR EQUAL PERIOD
CIRCULAR AND ELLIPTICAL ORBITS ABOUT NEPTUNE




Figure 19. Velocity change (ΔV) for Neptune.

ORIGINAL PAGE IS
OF POOR QUALITY

VELOCITY CHANGE VERSUS ORBIT ALTITUDE
FOR EQUAL PERIOD CIRCULAR AND ELLIPTICAL ORBITS ABOUT PLUTO





Figure 20. Velocity change (ΔV) for Pluto.

DEVELOPMENT OF EQUATIONS FOR EQUI-PERIOD ORBITS

The following assumptions were used in developing the equations for equi-period orbits (Fig 21):

- (a) A central force field.
- (b) A uniform homogeneous sphere of radius R_0 and acceleration of gravity g_0 .
- (c) A body M_1 in circular orbit at an altitude h above the surface of the sphere.
- (d) A body M_2 in an elliptical orbit with perigee altitude h_p and apogee altitude h_a above the surface of the sphere.



ORIGINAL PAGE IS
OF POOR QUALITY

Figure 21. Two-body orbital configuration.

To impose equi-period orbits, the time period for M_1 must equal the time period for M_2 :

$$\tau_{M_1} = \tau_{M_2} \quad (3)$$

The time period (τ_1) for M_1 in circular orbit is

$$\tau_{M_1} = \frac{2\pi \left(1 + \frac{h}{R_o}\right)^{3/2}}{\left(\frac{g_o}{R_o}\right)^{1/2}} \quad (4)$$

The time period for M_2 elliptical orbit is

$$\tau_{M_2} = \frac{2\pi}{\left(\frac{g_o}{R_o}\right)^{1/2}} \cdot \left(1 + \frac{h_p + h_a}{2R_o}\right)^{3/2} \quad (5)$$

Substituting equations (4) and (5) into equation (3) yields

$$\frac{2\pi \left(1 + \frac{h}{R_o}\right)^{3/2}}{\left(\frac{g_o}{R_o}\right)^{1/2}} = \frac{2\pi}{\left(\frac{g_o}{R_o}\right)^{1/2}} \left(1 + \frac{h_p + h_a}{2R_o}\right)^{3/2} \quad (6)$$

which simplifies to

$$\frac{h}{R_o} = \frac{h_p + h_a}{2R_o} \quad (7)$$


ORIGINAL PAGE IS
OF POOR QUALITY

and

$$h_a = 2h_a - h_p \quad . \quad (8)$$


We now select as our coordinate system a cartesian coordinate with origin at the center of the spheroid. The equation for an ellipse with one focal point at the origin of the coordinate system is

$$b^2 (x-c)^2 + a^2 y^2 = a^2 b^2 \quad . \quad (9)$$



The equation for a circle with center at the origin of the coordinate system is

$$x^2 + y^2 = r^2 \quad . \quad (10)$$



Rewriting equation (8),

$$h = \frac{1}{2} (h_a + h_p) \quad . \quad (11)$$

From Figure 21, it can be seen that the radius (r) of the circular orbit is equal to $h + R_o$. Adding R_o to both sides of equation (11),

$$r = h + R_o = \frac{1}{2} (h_a + h_p) + R_o \quad (12)$$

From Figure 21, the major axis $2a$ is

$$2a = h_p + 2R_o + h_a \quad (13)$$


The semimajor axis a is

$$a = \frac{1}{2} (h_p + h_a) + R_o \quad (14)$$

Comparing equations (12) and (14), it can be seen that the semimajor axis of the ellipse is equal to the radius of the circle in terms of R_o , h_a , and h_p .

As shown in Figure 21, the circle and ellipse intersect at coordinate $X = c$ and $Y = b$. From the trigonometric relationship of Figure 22,

$$a = R_o + h \quad (15)$$



ORIGINAL PAGE IS
OF POOR QUALITY

Figure 22. Orbital vectors.

In Figure 22,

dv = change of velocity required to transfer from a circular orbit of altitude h to an equi-period orbit of perigee h_p

R_o = radius of body

h = altitude of circular orbit

h_p = perigee of elliptical orbit

θ = change in direction of velocity from circular to equi-period elliptical orbit

V_e = elliptical velocity

V_c = circular velocity.

From the trigonometric relationship of Figure 21,

$$a - c - R_o - h_p = 0 \quad . \quad (16)$$

Substituting equation (15) into equation (16),

$$R_o + h - c - R_o - h_p = 0 \quad , \quad (17)$$

which simplifies to

$$c = h - h_p \quad . \quad (18)$$

From the trigonometric relationship of Figure 22,

$$b = (a^2 - c^2)^{1/2} \quad . \quad (19)$$

Substituting equations (15) and (17) into equation (19) yields

$$b = [(R_o + h)^2 - (h - h_p)^2]^{1/2} \quad . \quad (20)$$

From the trigonometric relationship of Figure 22, it can be seen that a is perpendicular to V_c and b is perpendicular to V_e . Therefore,

$$\theta = \alpha = \arctan \frac{c}{b} \quad . \quad (21)$$

Substituting equations (18) and (20) into equation (21) yields

$$\theta = \arctan \frac{h - h_p}{[(R_o + h)^2 - (h - h_p)^2]^{1/2}} \quad . \quad (22)$$

Dividing equation (22) by R,

$$\theta = \arctan \frac{\frac{h}{R_o} - \frac{h_p}{R_o}}{\left[\frac{(R_o + h)^2}{R_o^2} - \frac{(h - h_p)^2}{R_o^2} \right]^{1/2}} \quad . \quad (23)$$

Expanding the denominator,

ORIGINAL PAGE IS
OF POOR QUALITY

$$\theta = \arctan \frac{\frac{h}{R_o} - \frac{h_p}{R_o}}{\left[\frac{R_o^2}{R_o^2} + \frac{2 R_o h}{R_o^2} + \frac{h^2}{R_o^2} - \frac{h^2}{R_o^2} - \frac{2 h h_p}{R_o^2} - \frac{h_p^2}{R_o^2} \right]^{1/2}} , \quad (24)$$

which simplifies to

$$\theta = \arctan \frac{\frac{h}{R_o} - \frac{h_p}{R_o}}{\left[1 + \frac{2h}{R_o} + \frac{2h h_p}{R_o^2} - \frac{h_p^2}{R_o^2} \right]^{1/2}} . \quad (25)$$

From Figure 22, $V_e = V_c$ at points of intersection of circle and ellipse. Applying cosine law to Figure 22,

$$dv^2 = V_c^2 + V_c^2 - 2V_c^2 \cos \theta . \quad (26)$$

Taking square root of both sides,

$$dv = (2V_c^2 - 2V_c^2 \cos \theta)^{1/2} . \quad (27)$$

Simplifying,

$$dv = V_c (2 - 2 \cos \theta)^{1/2} . \quad (28)$$

From the trigonometric relationship of Figure 22,

$$\cos \theta = \cos \alpha = \frac{b}{a} ; \quad (29)$$

therefore,

$$dv = V_c \left(2 - \frac{2b}{a} \right)^{1/2} . \quad (30)$$

Substituting equation (15) for a and equation (20) for b into equation (30),

$$dv = V_c \left[\frac{2 - 2[(R_o + h)^2 - (h - h_p)^2]^{1/2}}{R_o + h} \right]^{1/2} . \quad (31)$$

Substituting $(g_o R_o / R_o + h)^{1/2}$ for V_c ,

$$dv = \left(\frac{g_o R_o}{R_o + h} \right)^{1/2} \left[2 - 2 \left[\frac{(R_o + h)^2 - (h - h_p)^2}{R_o + h} \right]^{1/2} \right]^{1/2} . \quad (30)$$

Collecting terms,

$$dv = R_o \sqrt{2 g_o} \left[\frac{1}{R_o + h} \left(1 - \left[\frac{(R_o + h)^2 - (h - h_p)^2}{R_o + h} \right]^{1/2} \right) \right]^{1/2} . \quad (31)$$

APPENDIX. VALUES USED FOR TABLES

	<u>Radius (km)</u>	<u>Acceleration of Gravity (m/sec/sec)</u>
Earth	6 371	9.806
Earth's Moon	1 738	1.622
Mercury	2 421	3.60
Venus	6 161	8.69
Mars	3 332	3.73
Jupiter	69 893	26.0
Saturn	57 532	11.2
Uranus	23 701	9.39
Neptune	21 535	14.99
Pluto	2 867	0.5

ORIGINAL PAGE IS
OF POOR QUALITY

BIBLIOGRAPHY

Escobol, P. R.: Methods of Orbit Determination. John Wiley, Inc., New York, 1965.


Herrick, Samuel and Baker, Robert: Gravitational and Related Constants for Accurate Space Navigation. The Eighth International Astronautical Congress, Barcelona, Spain, October 6, 1957.

APPROVAL

EQUAL PERIOD ORBIT DETERMINATION FOR THE SOLAR SYSTEM PLANETS

By Wilbur H. Funston

The information in this report has been reviewed for security classification. Review of any information concerning Department of Defense or nuclear energy activities or programs has been made by the MSFC Security Classification Officer. This report, in its entirety, has been determined to be unclassified.

A handwritten signature in cursive script, reading "J. T. Powell", is written over a horizontal line.

J. T. POWELL

Director, Data Systems Laboratory